


Di-electron Continuum at PHENIX

Y.L. Yamaguchi for the PHENIX collaboration

*Center for Nuclear Science, Graduate School of Science, University of Tokyo,
7-3-1 Hongo, Bunkyo, Tokyo, 113-0033, Japan*



The PHENIX experiment at RHIC has been carried out to investigate the properties of QGP. Di-electron yields in p+p and Au+Au collisions have been measured. An enhancement of the yield over a hadronic cocktail calculation is clearly seen for $p_T < 1.0$ GeV/ c and $150 < m_{ee} < 750$ MeV/ c^2 in Au+Au collisions, while the result in p+p collisions is consistent with the calculation. The fraction of the virtual direct photon component to the di-electron yield is measured from a shape analysis using the di-electron mass distributions for $1.0 < p_T < 5.0$ GeV/ c and $m_{ee} < 300$ MeV/ c^2 , and the real direct photon spectra are deduced from the fractions obtained for p+p and Au+Au collisions. An excess of the direct photon yield above a binary-scaled p+p result is seen in Au+Au collisions. The excess is fitted with an exponential function with an inverse slope parameter of $221 \pm 23 \pm 18$ MeV.

1 Introduction

The Quark Gluon Plasma (QGP) is a de-confined phase of a strongly-interacting matter. The lattice QCD calculation predicts a phase transition from a hadronic phase to QGP at a critical temperature of $150 \sim 200$ MeV. The study of QGP is very important since it gives not only new insights and intimate understanding of the strong interaction but also the understanding of the early universe at several μs after the Big Bang.

Heavy ion collision provides a unique tool to investigate the properties of QGP. A lot of very exciting results such as large energy loss of light quarks, gluons¹ and heavy quarks² and the strong elliptic flow for various particles^{2, 3} has been reported for $\sqrt{s_{NN}} = 200$ GeV Au+Au collisions at Relativistic Heavy Ion Collider (RHIC). These experimental facts indicate that the created matter has high-density and is rapidly-thermalized.

Di-leptons and direct photons are important probes to study QGP. They are emitted through all the stages of the collisions, and penetrate the strongly-interacting matter. Therefore they carry the thermodynamic information of the created matter directly. This paper focuses on

di-electron production in the low mass region below $1 \text{ GeV}/c^2$ and covers the following two topics. The first topic is the measurement of the di-electron yield in the low- p_T region, which is considered to be an optimal window for detecting thermal $q\bar{q}$ and $\pi\pi$ annihilations. The second one is the measurement of the direct photon yield with the virtual photon method ($\gamma^* \rightarrow e^+e^-$). Thermal photons from QGP are considered to be a primary contributor in $1.0 < p_T < 5.0 \text{ GeV}/c$ at RHIC energy⁴.

2 Low-mass Low- p_T Di-electron Continuum

The e^+e^- pair yield was measured in p+p and Au+Au collisions at PHENIX and hadron decay components were estimated using a hadronic cocktail calculation which incorporated the measured yields of the mesons for both p+p and Au+Au collisions^{5,6}. The left and right panels in Fig. 1 show the invariant mass spectra in p+p and Au+Au collisions compared to the hadronic cocktail calculations. The symbols and lines indicate the real data and the cocktail calculations, respectively. While the real data is in excellent agreement with the cocktail calculation in p+p collisions, a large enhancement over the calculation is clearly seen in $150 < m_{ee} < 750 \text{ MeV}/c^2$ in Au+Au collisions. The ratio of the integrated yield in $150 < m_{ee} < 750 \text{ MeV}/c^2$ between the real and the cocktail calculation is $3.4 \pm 0.2(\text{stat}) \pm 1.3(\text{sys}) \pm 0.7(\text{model})$. This enhancement is prominent in $p_T < 1.0 \text{ GeV}/c$ ⁷.




Figure 1: Invariant mass spectrum for p+p (left) and Au+Au (right) collisions compared to the hadronic cocktail calculations.

3 Low p_T Direct Photon

In general, any source of real photons can emit virtual photons which convert to low mass e^+e^- pairs. A direct photon production process such as $q + g \rightarrow g + \gamma$ has an associated process that produces low mass e^+e^- pairs, i.e. $q + g \rightarrow g + \gamma^* \rightarrow q + e^+e^-$. The relation between the photon production and the associated e^+e^- pair production is expressed by the Kroll-Wada formula, Eq. 1⁸.

$$\frac{d^2 n_{ee}}{dm_{ee}} = \frac{2\alpha}{3\pi} \frac{1}{m_{ee}} \sqrt{1 - \frac{4m_e^2}{m_{ee}^2}} \left(1 + \frac{2m_e^2}{m_{ee}^2} \right) S dn_\gamma, \quad (1)$$

where α is the fine structure constant, m_e and m_{ee} are the masses of the electron and the e^+e^- pair, respectively, and S is a process dependent factor.

In the case of hadrons such as π^0 and η Dalitz decays, S is given as $S = |F(m_{ee}^2)|^2 \left(1 - \frac{m_{ee}^2}{M_{hadron}^2} \right)^3$, where F denotes the form factor and M_{hadron} is the mass of the parent hadron. The S factor

is zero for $m_{ee} > M_{hadron}$. On the other hand, the S factor becomes unity for $m_{ee} \ll p_T$ in the case of virtual direct photon. Therefore it will be possible to extract the virtual direct photon component from the e^+e^- pair mass spectra by utilizing the difference in e^+e^- pair mass dependence of the S factor.




Figure 2: The left and center panels show the invariant mass spectra in p+p and Au+Au collisions for several p_T bins compared to the cocktail calculations. The right panel shows the mass spectrum in Au+Au collisions for $1.0 < p_T < 1.5$ GeV/c together with a fit result by Eq. 2.

An enhancement of the e^+e^- pair yield for Au+Au collisions is clearly visible in the p_T region above 1 GeV/c and the mass region of $100 < m_{ee} < 300$ MeV/c² as shown in the center panel of Fig. 2. We would like to emphasize that a small excess over the cocktail is also observed for p+p collisions in the high p_T region as shown in the left panel of Fig. 2.

The following function is used for fitting the data to determine the fraction of the virtual direct photon component in the mass spectrum.

$$f(m_{ee}) = (1 - r) \cdot f_{cocktail}(m_{ee}) + r \cdot f_{direct}(m_{ee}), \quad (2)$$

where $f_{cocktail}$ is the mass distribution from the decay of neutral hadrons estimated using the cocktail calculation and f_{direct} is that from the virtual direct photon decays, and r is the virtual direct photon fraction. Assuming that this excess comes from virtual direct photons, the fit to the result for Au+Au collisions in $1.0 < p_T < 1.5$ GeV/c gives $\chi^2/NDF = 13.8/10$. On the other hand, assuming that this excess comes from η , i.e. $f_\eta(m_{ee})$ is used in place of $f_{direct}(m_{ee})$, the fit gives $\chi^2/NDF = 21.1/10$. The result favors the virtual direct photons. It is also to be noted that ascribing this enhancement to η requires anomalous η enhancement, which seems very unlikely.

Figure 3 shows the obtained fractions of the virtual direct photon component as a function of p_T in p+p and Au+Au collisions. The symbols show the result and the lines are the expectations from next-to-leading-order perturbative QCD (NLO pQCD) calculations⁹ with different theoretical scales. A clear excess above the NLO pQCD calculation is seen in Au+Au collisions while the result in p+p collisions is consistent with the NLO pQCD calculation.

Finally, the real direct photon yield is obtained by multiplying the inclusive photon yield to the virtual direct photon fraction. Figure 4 shows the direct photon spectra in p+p and Au+Au collisions. This is the first time that the direct photon production in p+p collisions has been measured in $1.0 < p_T < 4.0$ GeV/c. The result in p+p collisions is well described by a modified power law function as shown by the dashed curve. The obtained yield in Au+Au collisions is above the N_{coll} -scaled p+p fitted curve in the low p_T region, where N_{coll} is the number of collisions. Fitting the result for Au+Au central collision with an exponential plus the N_{coll} -scaled p+p fitted curve provides the inverse slope parameter of $221 \pm 23 \pm 18$ MeV¹⁰. Comparing to available theoretical models gives a handle on the initial temperature that ranges from 300 to 600 MeV with $\tau_0 \sim 0.6-0.15$ fm/c. Implications of the results are under various investigations.




Figure 3: The obtained fractions of the virtual direct photon component as a function of p_T in p+p (left) and Au+Au (right) collisions.




Figure 4: The direct photon spectra in p+p and Au+Au collisions as a function of p_T .

References

1. K. Adcox *et al*, *Phys. Rev. Lett.* **88**, 022301 (2002).
2. K. Adcox *et al*, *Phys. Rev. Lett.* **98**, 172301 (2007).
3. S.S. Adler *et al*, *Phys. Rev. Lett.* **91**, 182301 (2003).
4. S. Turbide *et al*, *Phys. Rev. C* **69**, 014903 (2004).
5. A. Adare *et al*, arXiv:0802.0050 [nucl-ex].
6. S. Afanasiev *et al*, arXiv:0706.3034 [nucl-ex].
7. A. Toia, arXiv:0805.0153 [nucl-ex].
8. N.W. Kroll and W. Wada, *Phys. Rev.* **98**, 1355 (1955).
9. L.E. Gordon and W. Vogelsang, *Phys. Rev. D* **48**, 3136 (1993) and W. Vogelsang, private communication.
10. A. Adare *et al*, arXiv:0804.4168 [nucl-ex].